


Particle-decay branching ratios via intermediate-energy transfer reactions

Barry Davids


KVI, Groningen and
TRIUMF, Vancouver

Novae and x-ray bursts




R. Hynes 2000

Experimental Technique




- Populate resonances via surrogate reaction, study decays
- Studied $^{15}\text{O}(\alpha,\gamma)^{19}\text{Ne}$ using $p(^{21}\text{Ne},t)^{19}\text{Ne}$ reaction
- Excited states decay by α or γ emission
- Coincident detection of t ejectile and either ^{19}Ne recoil or ^{15}O decay product yields α -decay branching ratio


Experimental Setup at KVI's Big-Bite Spectrometer




Gamma Decay




Alpha Decay




Alpha Decays: Astrophysical Regime




^{22}Mg level scheme at energies relevant to $^{21}\text{Na}(p,\gamma)^{22}\text{Mg}$



Technique: proton-decay branching ratio measurement




Branching ratio data



Summary

- Measured α -decay branching ratios of states in ^{19}Ne and proton-decay branching ratios of states in ^{22}Mg
- Indirect technique requires independent lifetime measurements of particle-unbound states
- Allows determination of resonant reaction rates in small fraction of time required for direct measurements



Collaborators

J.P.M. Beijers, A.M. van den Berg, P.
Dendooven, S. Harmsma, M. Hunyadi,
M.A. de Huu, R.H. Siemssen, H.W.
Wilschut, and H.W. Wörtche

KVI, Groningen

K.E. Rehm and A.H. Wuosmaa

Argonne National Laboratory

R.E. Segel

Northwestern University

M. Hernanz and J. José

IEEC and UPC, Barcelona